

LETTER TO THE EDITOR

**A UV-crosslinkable interaction
in human U6 snRNA****JING-SONG SUN, SABA VALADKHAN, and JAMES L. MANLEY**Department of Biological Sciences, Sherman Fairchild Center of Life Science, Columbia University,
New York, New York 10027, USA**ABSTRACT**

U6 snRNA is the most conserved of all the snRNAs involved in pre-mRNA splicing, and likely plays an important role in splicing catalysis. Using a U6 snRNA fragment encompassing residues 25–99, we have identified a strong, UV-sensitive tertiary intramolecular interaction. A 5' deletion that removed sequences up to nt 37 only slightly reduced crosslinking, but further deletion of 11 bases, eliminating the nearly invariant ACAGAGA sequence, essentially abolished crosslinking, as did deletion of sequences 3' of 82A. The crosslinked residues were mapped to 44G in the ACAGAGA sequence and to 81C, the nucleotide at the base of the U6 intramolecular helix, opposite the G of the invariant AGC trinucleotide. This interaction is striking in that it has the potential to juxtapose invariant regions of U6 believed to play critical roles in splicing catalysis.

Keywords: pre-mRNA splicing; tertiary interaction; snRNA structure

Nuclear pre-mRNA splicing requires the assembly of the pre-mRNA substrate into the spliceosome, a dynamic structure consisting of a large number of distinct proteins as well as five small nuclear RNAs (snRNAs) (U1, U2, U4, U5, and U6). These snRNAs, which are preassembled into ribonucleoprotein particles (snRNPs), form extensive interactions with each other and with pre-mRNA. These interactions play important roles not only in spliceosome assembly, but also in the two catalytic transesterification steps in pre-mRNA splicing (reviewed by Madhani & Guthrie, 1994a; Nilsen, 1994; Sharp, 1994; Ares & Weiser, 1995). Four of the snRNAs participate in base pairing interactions with the pre-mRNA, although at most only three snRNAs, U2, U6, and U5, are likely to play direct roles in catalysis (Madhani & Guthrie, 1994a; Nilsen, 1994). This conclusion is based upon two types of experiments. First, biochemical assays, mainly UV and psoralen-induced crosslinking, have demonstrated that interactions between the pre-mRNA and all three of these snRNAs are formed within the spliceosome (Sawa & Abelson, 1992; Sawa & Shimura, 1992; Wassarman & Steitz, 1992; Wyatt

et al., 1992; Sontheimer & Steitz, 1993; Kim & Abelson, 1996). Second, genetic suppression studies have demonstrated that these interactions are functionally important (Parker et al., 1987; Wu & Manley, 1989; Zhuang & Weiner, 1989; Newman & Norman, 1991, 1992; Cortez et al., 1993; Kandels-Lewis & Seraphin, 1993; Lesser & Guthrie, 1993; Sun & Manley, 1995). However, the U5-pre mRNA interaction is dispensable for the first step of splicing *in vitro* (O'Keefe et al., 1996), and therefore only U2 and U6 are likely to play a direct role in catalysis, at least for the first step. These RNA–RNA interactions provide similarities to contacts that exist in autocatalytic group II introns (for review see Moore et al., 1993; Madhani & Guthrie, 1994a; see also Hetzer et al., 1997), and support the view that group II self-splicing and nuclear pre-mRNA splicing involve related mechanisms.

U6 snRNA is the most highly conserved of the snRNAs. Specifically, a central region of about 40 bases is nearly invariant between yeast and mammals (Brow & Guthrie, 1988; Fortner et al., 1994). Much of this conserved central domain is base paired with U4 snRNA in the U4/U6 snRNP (reviewed by Madhani & Guthrie, 1994a). U4/U6 and U5 snRNP are recruited to the spliceosome as a tri-snRNP complex. Upon completion of spliceosome assembly, the U4/U6 duplex dissoci-

Reprint requests to: James Manley, Department of Biological Sciences, Sherman Fairchild Center of Life Science, Columbia University, New York, New York 10027, USA; e-mail: jlm2@columbia.edu.

ates, and U4 likely leaves the spliceosome, allowing both extensive base pairing of U6 with U2 snRNA and formation of a U6 intramolecular helix. Both of these structures likely play important roles in splicing catalysis.

Genetic suppression experiments support the existence of three distinct U2-U6 helices, helix I, II, and III. Helix II was first identified by crosslinking and involves bases at the 5' end of U2 and 3' end of U6 (Hausner & Weiner, 1990), and the interaction was proven genetically in mammalian systems (Datta & Weiner, 1991; Wu & Manley, 1991). It may be of less significance in yeast, although a requirement for helix II was observed in a yeast strain carrying a mutation in U2-U6 helix Ib (see below), suggesting that these helices may be functionally redundant (Field & Friesen, 1996). Helix I was identified in yeast and was proposed to consist of two helices, Ia and Ib (Madhani & Guthrie, 1992, 1994b). This structure is striking in that bases involved in both U2 and U6 are adjacent to other critical bases. Helix Ia is situated just 5' to the nearly invariant ACAGAGA in U6, in which the underlined bases are implicated in 5' splice site pairing, and 3' to the U2 branch recognition sequence, which pairs with the pre-mRNA branch site. Helix Ia was proven subsequently in mammalian cells by genetic suppression (Sun & Manley, 1995), although current evidence suggests that helix Ib is not essential (Datta & Weiner, 1993; Sun & Manley, 1995). Finally, a third U2-U6 helix, helix III, has been proven by genetic suppression experiments in mammals (Sun & Manley, 1995). Although the potential to form helix III is well conserved evolutionarily (Takahashi et al., 1993), and it could further stabilize a 5' splice site-branch site interaction, it is not essential in yeast (Yan & Ares, 1996).

In addition to these intermolecular interactions, an intramolecular helix in U6 may also play an important role in splicing catalysis. Genetic experiments in both yeast (Fortner et al., 1994) and mammals (Wolff & Bindereif, 1993; Sun & Manley, 1995) have provided evidence that formation of the helix is necessary for some step(s) in splicing. Mutagenesis of human U6 revealed that the ability to participate in base pairing is compatible with function at all positions tested in this helix, and that bases not naturally paired could be incorporated into the helix without loss of function (Sun & Manley, 1997). These results suggest the role of the intramolecular helix may be largely structural and not sequence-specific.

Other studies suggest a similarity between the U6 intramolecular helix and domain 5 of group II introns. Analysis of critical phosphate oxygens in the two RNAs revealed that important phosphates were located at similar positions in secondary structure models of both RNAs (Fabrizio & Abelson, 1992; Chanfreau & Jaquier, 1994; Yu et al., 1995). Analysis of essential 2' hydroxyls also revealed similarities between the two structures (Abramovitz et al., 1996; Kim et al., 1997), and genetic analyses have suggested that corresponding

bases in U6 and domain 5 can be important for splicing (Madhani et al., 1990; Madhani & Guthrie, 1992; Boulanger et al., 1995; Peebles et al., 1995; Sun & Manley, 1995). U2-U6 helix Ib has been proposed to correspond to the bottom of domain 5 and the U6 intramolecular helix to the top, and genetic (Boulanger et al., 1995; Peebles et al., 1995) and chemical modification (Yu et al., 1995) studies are consistent with this model. Particularly important are the AGC bases at the bottom of both structures. These three bases are highly conserved in both systems and effects of mutations in both domain 5 and U6 are similar. However, a modification of this model, in which the U6 intramolecular helix is extended to replace helix Ib (Fortner et al., 1994), is more consistent with some mutational studies, at least in human U6 snRNA (Sun & Manley, 1995).

Here we describe a novel tertiary interaction, detected by UV crosslinking, in human U6 snRNA that bridges the invariant ACAGAGA sequence and the base of the U6 intramolecular helix. We show that crosslinking occurs between residue 44G, the central position in ACAGAGA, and 81C, which is at the base of the U6 intramolecular helix, opposite the G of the invariant AGC. This long-range tertiary interaction has the potential to juxtapose invariant regions of U6 that likely play critical roles in catalysis.

RESULTS

UV-induced crosslinking of human U6 snRNA

We undertook a UV crosslinking analysis of an in vitro-synthesized U6 snRNA fragment containing nucleotides 25–99 (U6 25–99). We chose this RNA because it includes the highly conserved central domain, the U6 sequences necessary for formation of U2-U6 helices I, II, and III, and the U6 intramolecular helix. In addition, bases 5' and 3' of this region have been shown to be unimportant for U6 function in vitro (Wolff & Bindereif, 1992). We were interested in the detection of possible interactions between and/or inside these functionally important domains, especially between ACAGAGA, helix I, and/or the U6 intramolecular helix. The U6 25–99 RNA was synthesized by in vitro transcription, dephosphorylated, and 5' end-labeled with [γ - P^{32}]ATP. In initial experiments, U6 25–99 was resuspended in a standard low-salt buffer used previously for group II self-splicing introns, containing 40 mM Tris-HCl, pH 7.5, 10 mM MgCl₂, and 2 mM spermidine (Peebles et al., 1987). The U6 RNA was first denatured at 70 °C for 2 min and then rapidly cooled on ice for 15 min. Renatured U6 snRNAs were UV irradiated and analyzed by denaturing gel electrophoresis. The heating and cooling procedure prior to irradiation was necessary for efficient crosslinking, presumably by allowing the RNA to assume a stably folded structure competent for crosslinking. Without heating and cooling, the same irradiation

procedure yielded less than 10% the amount of crosslinked RNA obtained with the renaturation procedure (results not shown).

Using the above protocol, a strong crosslinked product was detected in an initial experiment employing an RNA concentration of 100 nM. A time-course was performed to determine the efficiency of the crosslinking (Fig. 1). The crosslinked species formed with reasonable efficiency and accumulated up to 37% of the total RNA at 90 min, as determined by PhosphorImager analysis. The efficiency of crosslinking varied from about 20% to more than 50%. The lower yield likely reflects the fact that accumulation of crosslinked product did not always increase after 30–60 min, perhaps reflecting denaturation of the crosslinkable structure by excess heating during irradiation. The efficiency of crosslink formation indicates that, under these conditions, the crosslinkable conformation was a significant form of the U6 RNA fragment employed. This efficiency is comparable to those observed with several other RNAs (e.g., Branch et al., 1985, 1989; Behlen et al., 1992; Butcher & Burke, 1994), although less than with the

Tetrahymena group I ribozyme (Downs & Cech, 1990). To investigate whether the crosslink was intra- or intermolecular, crosslinking was performed in the presence of increasing concentrations of U625–99 RNA, from 0.1 nM to 10 μ M. The results (not shown) indicate that the rate and extent of crosslinking did not change significantly over the 100,000-fold variation in concentration, establishing that formation of the crosslink was independent of U6 concentration, and therefore likely intramolecular.

Sequence and ionic requirements for crosslinking

To investigate the regions of the U6 RNA fragment necessary for crosslinking, we examined several truncated derivatives of the U625–99 fragment. These include two 5' truncations beginning at positions 37 and 48, and a 3' truncation terminating at position 82. Note that the longer 5' truncation retains the ACAGAGA sequence, whereas the shorter begins just 3' to it. The 3' truncation terminates just one nucleotide 3' to the base of the intramolecular helix. RNAs were synthesized, dephosphorylated, 5' end-labeled, heated and cooled, and subjected to UV irradiation and denaturing PAGE as above. Figure 2 shows the results obtained after incubation for the indicated times, in buffer containing either 10 mM MgCl₂ or 1 mM EDTA. As above, U625–99 was crosslinked efficiently and in a Mg²⁺-dependent manner. Additional experiments (not shown) indicated that NaCl, over a concentration range of 50 mM to 1 M, could fully substitute for MgCl₂. These results suggest that MgCl₂ and NaCl function interchangeably to facilitate a structure compatible with crosslinking, likely by simple electrostatic interactions with the phosphate backbone. U637–99 also gave rise to a Mg²⁺-dependent crosslinked species of similar gel mobility, although the extent of crosslinking was reproducibly a factor of 2–3 lower than observed with U625–99. In contrast, crosslinking was undetectable with both U649–99 and U625–82 (results with U624–82 not shown). These results suggest that a residue or sequence in or adjacent to the ACAGAGA and 3' to the intramolecular helix is essential for crosslinking.

Identification of UV-crosslinked residues

To determine the identities of the crosslinked bases, both 5' and 3' end-labeled U625–99 RNAs were prepared, crosslinked under standard conditions (i.e., 40 mM Tris-HCl, pH 7.5, 10 mM MgCl₂, 2 mM spermidine), gel purified, and subjected to limited alkaline hydrolysis. The noncrosslinked RNAs were treated similarly, as well as subjected to partial digestion with a panel of RNases to provide a sequence ladder. Products were then resolved on 20% polyacrylamide gels, and results with the 5' (Fig. 3A) and 3' (Fig. 3B) labeled




FIGURE 1. Time-course of UV-induced crosslinking of U625–99 RNA. UV crosslinking was performed under standard conditions (top) for the indicated times, as described in Materials and Methods. RNAs were resolved by electrophoresis through an 8 M urea–20% polyacrylamide gel. XL and UNXL indicate the positions of crosslinked and uncrosslinked U6 RNAs, respectively. M is a DNA marker from 5'-labeled pBR322-*Hpa* II digestion.




FIGURE 2. RNA sequence requirements for crosslinking. Three U6 RNA fragments, U6 25–99, 37–99, and 48–99, were UV irradiated under the conditions indicated at the top of the panel (plus 40 mM Tris HCl, pH 7.5). Crosslinking was performed for the times indicated and RNAs resolved by denaturing gel electrophoresis as above. XL and UNXL indicate the positions of crosslinked and uncrosslinked U6 RNAs, respectively.

RNAs are shown. (Note that both of the purified cross-linked RNAs contained a small amount of uncross-linked RNA. Whether this was from contamination or from partial reversal of the crosslink is unclear, but its presence offers an explanation for a low background of hydrolysis products extending beyond the crosslinked base.) The 5'-labeled RNA revealed a distinct stop at residue A43, indicating that G44 is the major 5' cross-linked nucleotide. Samples in lanes XLOHa and XLOHb were made with different length of time of partial alkaline hydrolysis, 4.5 min and 5.5 min, respectively, and produced essentially identical results. Crosslinked species from NaCl-containing crosslinking buffer (40 mM Tris-HCl, pH 7.5, 100 mM NaCl) resulted in an identical stop at the same position (data not shown). With the 3' end-labeled RNA, a single stop at A82 was detected, implicating C81 as the 3' crosslinked nucleotide. Samples in lanes U6XL1a and 2a, U6XL1b and 2b were again alkaline-treated for 4.5 min and 5.5 min, respectively. Crosslinked products in lanes U6XL1a and b were made in the presence of NaCl, whereas cross-linked products in U6XL2a and b were from standard conditions. These data indicate that G44 and C81 participate in a long-range interaction within the U6 25–99 RNA molecule. Furthermore, these assignments are consistent with the results obtained with the truncation mutants described above. It is noteworthy that C81 is the penultimate 3' residue in the 3' truncation. Whether the failure of this RNA to crosslink reflects the terminal position of C81 in this RNA or a specific requirement for

downstream sequences is not known, but is discussed below.

DISCUSSION

Here we have described a specific UV-crosslinkable interaction between residues G44 and C81 in human U6 snRNA. Although most of our studies were performed with a truncated version of U6 snRNA, the same crosslink was detected with a full-length molecule (unpubl. data). This crosslink is similar to other tertiary RNA crosslinks described previously (e.g., Atmadja & Brimacombe, 1985; Branch et al., 1985, 1989; Downs & Cech, 1990; Butcher & Burke, 1994) in that it results from a folded structure of the U6 molecule, because it is detectable only after denaturing and annealing. The potential significance of this crosslink is enhanced because it juxtaposes G44 in the phylogenetically invariant ACAGAGA sequence and C81, which is opposite G54 in the invariant trinucleotide AGC. Thus, this interaction has the potential of bringing these two important regions of U6 snRNA into close proximity. Deletion mutations that remove either ACAGAGA or U6 sequences immediately downstream of C81 eliminated crosslinking, but single-nucleotide substitutions at either position unexpectedly had little effect (results not shown). These findings suggest that the overall structure of the region is more important for crosslinking than are the identities of the bases involved. We discuss below other




FIGURE 3. Identification of crosslinked residues in U6 25–99 RNA. The 5' (A) and 3' (B) crosslinked residues were mapped by limited alkaline hydrolysis (pH 12.0) of 5' and 3' end-labeled crosslinked and uncrosslinked products. U6 and U6XL represent purified uncrosslinked and crosslinked U6 25–99 RNA, respectively. U6 OH indicates partial alkaline hydrolysis of the uncrosslinked RNA. U6 XL a and b were subjected to alkaline hydrolysis for 4.5 and 5.5 min, respectively. In B, XL1 depicts RNA crosslinked in the presence of 100 mM NaCl, whereas XL2 is from reactions performed under standard conditions (see text). 5'- or 3'-Labeled uncrosslinked U6 RNAs were partially digested with RNases T1 (G), U2 (A), C13 (C), and PhyM (A+U) to produce a sequence ladder. RNAs were resolved by denaturing 20% PAGE. Crosslinked residues are indicated with arrows.

potential interactions that may contribute to formation of the crosslinkable configuration.

The tertiary interaction involving G44 and C81 is interesting in that previous studies have suggested potentially important roles for these bases in splicing, especially for G44. For example, the single-base substitution G44C has been shown to abolish splicing activity in both transient expression in vivo (Datta & Weiner, 1993) and splicing complementation in vitro (Wolff et al., 1994). A G → C mutation at the corresponding position in yeast U6 snRNA led to a significant reduction in in vitro reconstitution activity (Fabrizio & Abelson, 1990), and to a *ts* growth phenotype in vivo (Madhani et al., 1990; McPheeters, 1996). Both G44U and G44A mutations can be lethal in yeast (McPheeters, 1996), although G44A had no effect on splicing in transfected human cells (Datta & Weiner, 1993). Mutation of C81 to G resulted in a decrease in splicing in a transient expression assay, although the defect could be rescued by a compensatory G → C change in the essential residue G54 (Sun & Manley, 1995). A number of previous studies have illustrated the importance of G54, not only in yeast (e.g., McPheeters, 1996) and mammalian (Datta & Weiner, 1993; Sun & Manley, 1995) splicing, but also in group II splicing, where a comparable AGC sequence is situated at the base of domain 5 (Boulanger et al., 1995; Peebles et al., 1995).

A number of additional studies have indicated the importance for splicing of bases and/or phosphate oxygens near the two positions identified in our crosslinking studies. Crosslinking experiments in yeast and humans have shown that the ACA of the ACAGAGA sequence physically contacts the pre-mRNA 5' splice site, an interaction that has also been proven genetically in yeast and mammals (see Introduction for references). Thiophosphate incorporation at various positions in U6 RNA has identified important phosphate oxygen atoms in both yeast and nematodes (Fabrizio & Abelson, 1992; Yu et al., 1995). The phosphate oxygen between ACA and GAGA was shown to be critical for splicing activity in both organisms, which may be particularly significant in light of our crosslinking studies implicating this same position in the intramolecular crosslink. The phosphate oxygen immediately 5' to the invariant AGC was shown to be necessary for the second step of splicing (Fabrizio & Abelson, 1992; Yu et al., 1995), whereas a thiophosphate 5' to the G inhibited splicing in worms (Yu et al., 1995) and also in group II self splicing (Chanfreau & Jacquier, 1994). U6-81C is in close proximity to these two oxygens via base pairing with G54 (Sun & Manley, 1995), and the crosslinkable conformation we have described thus has the potential to help bring all three of these important phosphate oxygens into close proximity. An intriguing possibility is that this contributes to formation of a metal binding site essential for catalysis (e.g., Steitz & Steitz, 1993).

Several related secondary structures for free human U6 snRNA have been proposed (Epstein et al., 1980; Harada et al., 1980; Miura et al., 1983; Rinke et al., 1985). The structure illustrated in Figure 4A is similar but not identical to the earlier structures. These structures all share in common what is now known as the U6 intramolecular helix, which extends from G54 to C81. Most of the paired positions have been proven to exist at some stage during splicing by genetic suppression experiments (Wolff & Bindereif, 1993; Sun & Manley, 1995, 1997). The proposed structures also all share the lower stem, although we are not aware of direct evidence, genetic or chemical, confirming its existence or importance during splicing. The central loop and helix shown in Figure 4A have been drawn in several configurations. There is again limited evidence for or against any of these structures, although chemical modification data suggest that C42 is unpaired (Miura et al., 1983).

The secondary structure depicted in Figure 4A has a striking feature not noticed previously, which is a remarkable similarity with the currently envisioned U2-U6 secondary structure, excluding U2-U6 helix II (Fig. 4B). It shows not only structural similarity, but also nucleotide identities at several functionally important positions, such as U2-U6 helix Ia and a region overlapping the U2 branch site recognition sequence, AUGAUG (identities are lightly shaded in Fig. 4). Whether these similarities are functionally significant is unclear, but they do raise the possibility that the U6 RNA fragment we have analyzed reflects the U2-U6 structure thought to exist in the spliceosome. In addition, this structure provides a framework to consider the basis for the effects of the deletions on crosslinking. Removal of bases 5' to 37C reduced crosslinking efficiency, and eliminated four of six bases from the bottom stem, consistent with the possibility that this helix may contribute to formation of the crosslinkable conformation. Deletion of bases 3' to 82A, which eliminated crosslinking, would disrupt all the structure below the intramolecular helix. Future experiments will be aimed at determining in more detail the role of the proposed U6 structure in crosslinking, as well as whether the 44G-81C crosslinkable conformation exists in the spliceosome, when U6 is paired to U2.

By using an elegant in vivo bimolecular randomization-selection assay, Madhani and Guthrie (1994b) obtained genetic evidence in yeast for a tertiary interaction between the penultimate nucleotide in the ACAGAGA sequence and the 3' base in the two-nucleotide bulge between U2-U6 helices Ia and Ib (Fig. 4C). This interaction has been suggested to contribute to formation of the catalytic center for the second step of splicing by helping to juxtapose the ACAGAGA and AGC sequences. The similarity between this interaction and the one described here is striking, and it is conceivable that both could exist simultaneously in the same folded




FIGURE 4. Potential U6 and U2-U6 snRNA secondary structures in human and yeast. Numbers indicate nucleotide positions from 5' end. Dark shading highlights the invariant ACAGAGA and AGC sequences. Light shading denotes sequence identities between U6 RNA and human and yeast U2 RNA in the regions corresponding to helix Ia and the branch site recognition region (see text). **A:** U6 RNA secondary structure model, residues 33–99. Crosslinked residues are indicated by asterisks and joined by dashed arrow. **B:** Human U6-U2 structure, including helices Ia and III and the intramolecular helix. **C:** Yeast U6-U2 structure, including helices Ia and b, putative helix III, and the intramolecular helix. Residues involved in a genetically established tertiary interaction (Madhani & Guthrie 1994b) are indicated.

RNA structure. There are minor differences between the relevant regions in the proposed yeast and human secondary structures (e.g., yeast helix Ib is replaced by the extended U6 intramolecular helix in human; compare Fig. 4B and C), but these may not be functionally important, because both structures can form related secondary and, potentially, tertiary structures. However, in yeast, a U2 snRNA residue, 22C, would be predicted to play the role of U681C in human. Or, given the lack of importance of base identity at the crosslinkable positions, perhaps the human equivalent of the yeast interaction, which would involve U6 residues 46G and 82A, plays a key role in establishing the conformation that gives rise to the crosslink we have analyzed. In any event, it will be important in the future to determine how the tertiary interaction we have described contributes to the functions of U6 snRNA in splicing.

MATERIALS AND METHODS

RNA transcription and end-labeling

All U6 snRNA fragments were transcribed from DNA templates that were amplified by PCR on a U6-containing plasmid by a 5'-overhanging primer containing the sequence of the T7 promoter region and transcriptions were performed with bacteriophage T7 RNA polymerase. All RNAs were purified using 20% denaturing PAGE in Tris-borate-EDTA buffer, eluted by overnight diffusion, and precipitated. RNAs were 5'-labeled with [γ - 32 P]ATP and T4 polynucleotide kinase following dephosphorylation with calf intestinal phosphatase, or 3' end-labeled with [5'- 32 P]pCp and T4 RNA ligase. Heterogeneity resulting from the addition of nontemplated nucleotides by T7 RNA polymerase could be eliminated in the case of the 3' end-labeled RNAs by long runs on 6% denaturing polyacrylamide gels (80-cm plate) at 1,600 V for 8 h.

UV crosslinking

After end-labeling and gel purification, the appropriate U6 RNA species were resuspended in 40 mM Tris-HCl, pH 7.5, and 10 mM MgCl₂ unless otherwise noted. Unlabeled U6 RNAs were added to adjust the final concentration to 100 nM, unless otherwise noted. The RNAs were allowed to fold prior to UV irradiation by heating to 70 °C for 2 min and then rapidly cooling on ice for 15 min. Heating the RNA to any temperature above 60 °C and cooling either quickly or gradually gave the same extent of crosslinking (results not shown). Twenty-microliter aliquots were placed into wells of a Falcon 96-well microtiter plate resting on ice. Samples were irradiated for 10-min intervals interrupted by 10 min to prevent overheating. Irradiation was done with a hand-held 254-nm UV light (model UVG-54, UV Products Inc.; measured output, 2,200 μW/cm²) at a distance of 1 cm. After the indicated times (0–90 min), RNAs were directly electrophoresed on 20% denaturing polyacrylamide gels. For mapping of the crosslinks, RNA products were visualized by autoradiography, excised from the gel, eluted by diffusion in 0.3 M ammonium acetate overnight, and precipitated with 2.5 volumes of ethanol. Quantitation of crosslinking was done by using a Storm PhosphorImager (Molecular Dynamics).

Mapping of crosslinked bases

To map the crosslinks, gel-purified, end-labeled crosslinked RNAs were subjected to partial alkaline hydrolysis. Samples were dissolved in 50 mM (NH₄)₃PO₄ buffer, pH 12.0, heated to 100 °C for 4.5 or 5.5 min for hydrolysis of crosslinked RNA, and for 30 s for hydrolysis of uncrosslinked RNA. The partially hydrolyzed samples were loaded directly onto a 20% denaturing polyacrylamide gel. Noncrosslinked RNAs were partially digested with base-specific ribonucleases T1, U2, C13, and PhyM (U.S. Biochemical) to provide sequencing ladders to identify the positions of the crosslinked bases.

ACKNOWLEDGMENTS

We thank Yoshio Takagaki, Shou-Hua Xiao, Jin Wang, and Chi Li for their discussion and generous suggestions; Lin Ge and Xiaohong Shi for excellent technical assistance; and John Burke and Joyce Heckman for kindly providing in vitro transcription template for production of the 51-nt hairpin ribozyme, and for their experimental protocols for partial alkaline hydrolysis and partial RNase digestion. This work was supported by NIH grant GM 48259.

Received January 26, 1998; returned for revision February 16, 1998; revised manuscript received February 27, 1998

REFERENCES

- Abramovitz DL, Friedman RA, Pyle AM. 1996. Catalytic role of 2'-hydroxyl groups within a group II intron active site. *Science* 271:1410–1413.
- Ares M Jr, Weiser B. 1995. Rearrangement of snRNA structure during assembly and function of the spliceosome. *Prog Nucleic Acid Res Mol Biol* 50:131–159.
- Atmadja J, Brimacombe R, Blocker H, Frank R. 1985. Investigation of the tertiary folding of *Escherichia coli* 16S RNA by in situ intramolecular RNA cross-linking within 30S ribosomal subunits. *Nucleic Acids Res* 13:6919–6936.
- Behlen LS, Sampson JR, Uhlenbeck OC. 1992. An ultraviolet light-induced crosslink in yeast tRNA(Phe). *Nucleic Acids Res* 20:4055–4059.
- Boulanger SC, Belcher SM, Schmidt U, Dib-Hajj SD, Schmidt T, Perlman PS. 1995. Studies of point mutants define three essential paired nucleotides in the domain 5 substructure of a group II intron. *Mol Cell Biol* 15:4479–4488.
- Branch AD, Benenfeld BJ, Baroudy BM, Wells FV, Gerin JL, Robertson HD. 1989. An ultraviolet-sensitive RNA structural element in a viroid-like domain of the hepatitis delta virus. *Science* 243:649–652.
- Branch AD, Benenfeld BJ, Robertson HD. 1985. Ultraviolet light-induced crosslinking reveals a unique region of local tertiary structure in potato spindle tuber viroid and HeLa 5S RNA. *Proc Natl Acad Sci USA* 82:6590–6594.
- Brow DA, Guthrie C. 1988. Spliceosomal RNA U6 is remarkably conserved from yeast to mammals. *Nature* 334:213–218.
- Butcher SE, Burke JM. 1994. A photo-cross-linkable tertiary structure motif found in functionally distinct RNA molecules is essential for catalytic function of the hairpin ribozyme. *Biochemistry* 33:992–999.
- Chanfreau G, Jacquier A. 1994. Catalytic site components common to both splicing steps of a group II intron. *Science* 266:1383–1387.
- Cortez JJ, Sontheimer EJ, Seiwert SD, Steitz JA. 1993. Mutations in the conserved loop of human U5 snRNA generate use of novel cryptic 5' splice sites in vivo. *EMBO J* 12:5181–5189.
- Datta B, Weiner AM. 1991. Genetic evidence for base pairing between U2 and U6 snRNAs in mammalian mRNA splicing. *Nature* 352:821–824.
- Datta B, Weiner AM. 1993. The phylogenetically invariant ACAGAGA and AGC sequences of U6 small nuclear RNA are more tolerant of mutation in human cells than in *Saccharomyces cerevisiae*. *Mol Cell Biol* 13:5377–5382.
- Downs WD, Cech TR. 1990. An ultraviolet-inducible adenosine-adenosine cross-link reflects the catalytic structure of the *Tetrahymena* ribozyme. *Biochemistry* 29:5605–5613.
- Epstein P, Reddy R, Henning D, Busch H. 1980. The nucleotide sequence of nuclear U6 (4.7 S) RNA. *J Biol Chem* 255:8901–8906.
- Fabrizio P, Abelson J. 1990. Two domains of yeast U6 small nuclear RNA required for both steps of nuclear precursor messenger RNA splicing. *Science* 250:404–409.
- Fabrizio P, Abelson J. 1992. Thiophosphates in yeast U6 snRNA specifically affect pre-mRNA splicing in vitro. *Nucleic Acids Res* 20:3659–3664.
- Field DJ, Friesen JD. 1996. Functionally redundant interactions between U2 and U6 spliceosomal snRNAs. *Genes & Dev* 10:489–501.
- Fortner DM, Troy RG, Brow DA. 1994. A stem/loop in U6 RNA defines a conformational switch required for pre-mRNA splicing. *Genes & Dev* 8:221–233.
- Harada F, Kato N, Nishimura S. 1980. The nucleotide sequence of nuclear 4.8S RNA of mouse cells. *Biochem Biophys Res Commun* 95:1332–1340.
- Hausner TP, Giglio LM, Weiner AM. 1990. Evidence for base-pairing between mammalian U2 and U6 small nuclear ribonucleoprotein particles. *Genes & Dev* 4:2146–2156.
- Hetzer M, Wurzer G, Schweyen RJ, Mueller MW. 1997. Trans-activation of group II intron splicing by nuclear U5 snRNA. *Nature* 386:417–420.
- Kandels-Lewis S, Seraphin B. 1993. Role of U6 snRNA in 5' splice site selection. *Science* 262:2035–2039.
- Kim CH, Abelson J. 1996. Site-specific crosslinks of yeast U6 snRNA to the pre-mRNA near the 5' splice site. *RNA* 2:995–1010.
- Kim CH, Ryan DE, Marciniak J, Abelson J. 1997. Site-specific deoxynucleotide substitutions in yeast U6 snRNA block splicing of pre-mRNA in vitro. *EMBO J* 16:2119–2129.
- Lesser CF, Guthrie C. 1993. Mutations in U6 snRNA that alter splicing site specificity: Implications for the active site. *Science* 262:1982–1988.
- Madhani HD, Bordonne R, Guthrie C. 1990. Multiple roles for U6 snRNA in the splicing pathway. *Genes & Dev* 4:2264–2277.

- Madhani HD, Guthrie C. 1992. A novel base-pairing interaction between U2 and U6 snRNAs suggests a mechanism for the catalytic activation of the spliceosome. *Cell* 71:803–817.
- Madhani HD, Guthrie C. 1994a. Dynamic RNA–RNA interactions in the spliceosome. *Annu Rev Genet* 28:1–26.
- Madhani HD, Guthrie C. 1994b. Randomization-selection analysis of snRNAs in vivo: Evidence for a tertiary interaction in the spliceosome. *Genes & Dev* 8:1071–1086.
- McPheeters DS. 1996. Interactions of the yeast U6 RNA with the pre-mRNA branch site. *RNA* 2:1110–1123.
- McPheeters DS, Abelson J. 1992. Mutational analysis of the yeast U2 snRNA suggests a structural similarity to the catalytic core of group I introns. *Cell* 71:819–831.
- Miura K, Tsuda S, Harada F, Ueda T. 1983. Chemical modification of cytosine residues of U6 snRNA with hydrogen sulfide. *Nucleic Acids Res* 11:5893–5901.
- Moore MJ, Query CC, Sharp PA. 1993. Splicing of precursors to mRNA by the spliceosome. In: Gesteland RF, Atkins JF, eds. *The RNA world*. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press. pp 303–358.
- Newman AJ, Norman C. 1991. Mutations in yeast U5 snRNA alter the specificity of 5' splicing-site cleavage. *Cell* 65:115–123.
- Newman AJ, Norman C. 1992. U5 snRNA interacts with exon sequences at 5' and 3' splice sites. *Cell* 68:743–754.
- Nilsen TW. 1994. RNA–RNA interactions in the spliceosome: Unraveling the ties that bind. *Cell* 78:1–4.
- O'Keefe RT, Newman C, Norman AJ. 1996. The invariant U5 snRNA loop 1 sequence is dispensable for the first catalytic step of pre-mRNA splicing in yeast. *Cell* 86:679–689.
- Parker R, Siliciano PG, Guthrie C. 1987. Recognition of the TAC TAAC box during mRNA splicing in yeast involves base pairing to the U2-like snRNA. *Cell* 49:229–239.
- Peebles CL, Benatan EJ, Jarrell KA, Perlman PS. 1987. Group II intron self-splicing: Development of alternative reaction conditions and identification of a predicted intermediate. *Cold Spring Harbor Symp Quant Biol* 52:223.
- Peebles CL, Zhang M, Perlman PS, Franzen JS. 1995. Catalytically critical nucleotides in domain 5 of a group II intron. *Proc Natl Acad Sci USA* 92:4422–4426.
- Rinke J, Appel B, Digweed M, Luhrmann R. 1985. Localization of a base-paired interaction between small nuclear RNAs U4 and U6 in intact U4/U6 ribonucleoprotein particles by psoralen cross-linking. *J Mol Biol* 185:721–731.
- Sawa H, Abelson J. 1992. Evidence for a base-pairing interaction between U6 small nuclear RNA and 5' splice site during the splicing reaction in yeast. *Proc Natl Acad Sci USA* 89:11269–11273.
- Sawa H, Shimura Y. 1992. Association of U6 snRNA with the 5'-splice site region of pre-mRNA in the spliceosome. *Genes & Dev* 6:244–254.
- Sharp PA. 1994. Split genes and RNA splicing. *Cell* 77:805–815.
- Sontheimer E, Steitz JA. 1993. The U5 and U6 small nuclear RNAs as active site components of the spliceosome. *Science* 262:1989–1996.
- Steitz TA, Steitz JA. 1993. A general two-metal-ion mechanism for catalytic RNA. *Proc Natl Acad Sci USA* 90:6498–6502.
- Sun JS, Manley JL. 1995. A novel U2-U6 snRNA structure is necessary for mammalian mRNA splicing. *Genes & Dev* 9:843–854.
- Sun JS, Manley JL. 1997. The human U6 snRNA intramolecular helix: Structural constraints and lack of sequence specificity. *RNA* 3:514–526.
- Takahashi Y, Urushiyama S, Tani T, Ohshima Y. 1993. An mRNA-type intron is present in the *Rhodotorula hasegawae* U2 small nuclear RNA gene. *Mol Cell Biol* 13:5613–5619.
- Wassarman DA, Steitz JA. 1992. Interactions of small nuclear RNA's with precursor messenger RNA during in vitro splicing. *Science* 257:1918–1925.
- Wolff T, Bindereif A. 1992. Reconstituted mammalian U4/U6 snRNP complements splicing: A mutational analysis. *EMBO J* 11:345–359.
- Wolff T, Bindereif A. 1993. Conformational changes of U6 RNA during the spliceosomal cycle: An intramolecular helix is essential both for initiating the U4-U6 interaction and for the first step of splicing. *Genes & Dev* 7:1377–1389.
- Wolff T, Menssen R, Hammel J, Bindereif A. 1994. Splicing function of mammalian U6 small nuclear RNA: Conserved positions in central domain and helix I are essential during the first and second step of pre-mRNA splicing. *Proc Natl Acad Sci USA* 91:903–907.
- Wu J, Manley JL. 1989. Mammalian pre-mRNA branchsite selection by U2 snRNP involves base pairing. *Genes & Dev* 3:1553–1561.
- Wu J, Manley JL. 1991. Base-pairing between U2 and U6 snRNAs is necessary for splicing of a mammalian pre-mRNA. *Nature* 352:818–821.
- Wyatt JR, Sontheimer EJ, Steitz JA. 1992. Site-specific cross-linking of mammalian U5 snRNP to the 5' splice site before the first step of pre-mRNA splicing. *Genes & Dev* 6:2542–2553.
- Yan D, Ares M. 1996. Invariant U2 RNA sequences bordering the branchpoint recognition region are essential for interaction with yeast SF3a and SF3b subunits. *Mol Cell Biol* 16:818–828.
- Yu YT, Maroney PA, Darzynkiwicz E, Nilsen TW. 1995. U6 snRNA function in nuclear pre-mRNA splicing: A phosphorothioate interference analysis of the U6 phosphate backbone. *RNA* 1:46–54.